

Growth mixture models in longitudinal research

Jost Reinecke · Daniel Seddig

Received: 20 August 2010 / Accepted: 3 March 2011 / Published online: 19 October 2011
© Springer-Verlag 2011

Abstract Latent growth curve models as structural equation models are extensively discussed in various research fields (Curran and Muthén in Am. J. Community Psychol. 27:567–595, 1999; Duncan et al. in An introduction to latent variable growth curve modeling. Concepts, issues and applications, 2nd edn., Lawrence Erlbaum, Mahwah, 2006; Muthén and Muthén in Alcohol. Clin. Exp. Res. 24(6):882–891, 2000a; in J. Stud. Alcohol. 61:290–300, 2000b). Recent methodological and statistical extension are focused on the consideration of unobserved heterogeneity in empirical data. Muthén extended the classic structural equation approach by mixture components, i.e. categorical latent classes (Muthén in Marcoulides, G.A., Schumacker, R.E. (eds.), New developments and techniques in structural equation modeling, pp. 1–33, Lawrence Erlbaum, Mahwah, 2001a; in Behaviometrika 29(1):81–117, 2002; in Kaplan, D. (ed.), The SAGE handbook of quantitative methodology for the social sciences, pp. 345–368, Sage, Thousand Oaks, 2004). The paper discusses applications of growth mixture models with data on delinquent behavior of adolescents from the German panel study *Crime in the modern City (CrinoC)* (Boers et al. in Eur. J. Criminol. 7:499–520, 2010; Reinecke in Delinquenzverläufe im Jugendalter: Empirische Überprüfung von Wachstums- und Mischverteilungsmodellen, Institut für sozialwissenschaftliche Forschung e.V., Münster, 2006a; in Methodology 2:100–112, 2006b; in van Montfort, K., Oud, J., Satorra, A. (eds.), Longitudinal models in the behavioral and related sciences, pp. 239–266, Lawrence Erlbaum, Mahwah, 2007). Observed as well as unobserved heterogeneity will be considered with growth mixture models. Special attention is given to the distribution of the outcome variables as

J. Reinecke (✉)

Faculty of Sociology, University of Bielefeld, Bielefeld, Germany
e-mail: jost.reinecke@uni-bielefeld.de

D. Seddig

Department of Criminology, University of Münster, Münster, Germany
e-mail: d.seddig@uni-muenster.de

counts. Poisson and negative binomial distributions with zero inflation are considered in the proposed growth mixture models variables. Different model specifications will be emphasized with respect to their particular parameterizations.

Keywords Panel data · Latent class growth analysis · Growth mixture modeling · Heterogeneity · Zero-inflated negative binomial model

1 Introduction

Longitudinal research studies with repeated measurements are quite often used to examine processes of stability and change in individuals or groups. With panel data it is possible to investigate intraindividual development of substantive variables across time as well as interindividual differences and similarities in change patterns. While the traditional analysis of variance (ANOVA) and the analysis of covariance (ANCOVA) assume homogeneity of the underlying covariance matrix across the levels of the between-subjects factors and the same covariance patterns for the repeated measurements, the structural equation methodology offers an alternative strategy: the *latent growth curve models*. These models describe not only a single individual's developmental trajectory, but also capture individual differences in the intercept and slopes of those trajectories. Based on the formative work of Rao and Tucker's basic model of growth curves (Rao 1958; Tucker 1958), Meredith and Tisak (1990) discussed and formalized the model within the structural equation framework. Further developments of the growth curve model were proposed by McArdle and Epstein (1987), McArdle (1988) and Muthén (1991, 1997).

Observed heterogeneity in growth curve models can be captured by covariates explaining part of the variances of the intercept and slope. But the assumption of a single population underlying the growth curves has to be relaxed in the case of unobserved heterogeneity. Instead of considering individual variation around a single growth curve, different classes of individuals should vary around different mean growth curves. A very suitable framework to handle the issue of unobserved heterogeneity is *growth mixture modeling* introduced by Muthén and Shedden (1999). These mixture models differ between continuous and categorical latent variables. The categorical latent variables represent mixtures of subpopulations where the product membership is inferred from the data. Like the conventional growth curve models, intercept and slope variables capture the continuous part of the model. Growth mixture models can also be seen as an extension of the structural modeling approach with techniques of latent class analysis. The inferred membership of each individual to a certain class is produced with the information of the estimated latent class probabilities. Further developments and applications with the program *Mplus* (Muthén and Muthén 2006) are discussed in several papers by Muthén (2001a, 2001b, 2003, 2004). Recently, Muthén (2008) gives a model overview of the so-called *latent variable hybrids* within the continuous and categorical latent variable framework.

Statistical techniques to group and classify individuals with categorical panel data (e.g., latent class analysis) have a long tradition in behavioral and social sciences (Rost and Langeheine 1997). In criminology, classification of longitudinal continuous and count data were formally introduced by Nagin and Land (1993), Nagin

(1999) and Roeder, Lynch and Nagin (1999) with the *semiparametric group-based approach* (see also Nagin 2005 for an overview). This technique enhanced the ability to estimate group membership of individuals who follow a common trajectory across time (e.g., persistent offenders). Muthén (2004) discusses this group-based approach as a simpler specification of the general growth mixture model and labeled it as *latent class growth analysis*. In difference to the general growth mixture model the growth curve parameters are fixed and not random, assuming no variation across individuals within classes. The possibility to treat their measurements as counts with the Poisson distribution as the underlying statistical model (see, e.g., Ross 1993) is also part of the mixture model. If the count variables are inflated with zeros, i.e. the particular behaviors seldom occur, a variant of the Poisson model, the so-called zero-inflated Poisson model (ZIP, Lambert 1992) should lead to a better statistical representation of the data than a model without considering the zero inflation. However, a Poisson distribution assumes the equality of its mean and variance. This property is rarely found in empirical data. If the variance is larger than the mean, then the negative binomial model (NB) should be used instead of the Poisson model to get a parameter estimate of the overdispersion (cf. Hilbe 2007 for an overview of the varieties of negative binomial models). If the count variables are both inflated with zeros and overdispersed, the so-called zero-inflated negative binomial model (ZINB, Hilbe 2007, p. 174f.) should be applied. The recent version of the *Mplus* program allows the specification of different count data models including the ZIP- and the ZINB model (Muthén and Muthén 2010). The paper will focus on the applicability of zero-inflated count data models which has been seldom used within growth mixture analyses.

After a short introduction of growth curve and growth mixture models, special cases for count data are discussed in Section 2. Section 3 gives a brief introduction of the longitudinal study, the sample and the variables used in the statistical analyses. Results of the growth curve and growth mixture models are presented in Sections 4 and 5. A summary and discussion with suggestions for further research are given in Section 6.

2 Method and models


2.1 Growth curve models

The possibility that the individual trajectories of a dependent variable can vary is one of the main advantages of the growth curve model. The formal representation of a growth curve model can be seen either as a multilevel, random-effects model or as a latent variable model, where the random effects are latent variables (Meredith and Tisak 1990, p. 108; Willet and Sayer 1994, p. 369):

$$y_i = \Lambda \eta_i + \epsilon_i \quad (1)$$

y_i is a $t \times 1$ vector of repeated measurements for observation i where t is the number of panel waves. η is a $q \times 1$ vector of latent growth factors where q is the number of these factors. ϵ is a $t \times 1$ vector of time-specific measurement errors, and Λ is the $t \times q$ matrix of factor loadings with fixed coefficients representing the functional form

Fig. 1 Quadratic growth curve model for t panel waves



of the individual trajectories. Variations of individual trajectories are captured by q -numbers of latent variables η whereas usually η_1 is the *intercept*, η_2 is the *linear slope* and in case of nonlinear development η_3 represents the *quadratic slope* (cf. Fig. 1). If applicable, additional latent variables can be specified. It is assumed that the latent growth factors and measurement errors are independent and multivariate normally distributed:

$$\begin{bmatrix} \eta_i \\ \epsilon_i \end{bmatrix} \approx \left(\begin{bmatrix} \alpha \\ 0 \end{bmatrix}, \begin{bmatrix} \Psi & 0 \\ 0 & \Theta \end{bmatrix} \right) \quad (2)$$

where α is a $q \times 1$ vector of growth factor means and Ψ is the respective $q \times q$ covariance matrix. Θ is a $p \times p$ covariance matrix of time-specific measurement errors which are usually constrained to be a diagonal matrix.

For estimation a probability density function is used:

$$f(y_i) = \phi[y_i; \mu(\theta) \Sigma(\theta)] \quad (3)$$

where ϕ is the probability density function for y_i and θ is the vector of all parameters to be estimated. $\mu(\theta)$ is a $p \times 1$ model-implied mean vector given by

$$\mu(\theta) = \Lambda \alpha \quad (4)$$

and $\Sigma(\theta)$ is a $p \times p$ model-implied covariance matrix given by

$$\Sigma(\theta) = \Lambda \Psi \Lambda' + \Theta. \quad (5)$$

Parameters in θ are ML estimates which maximizes the likelihood that the measurements y_i are drawn from a multivariate normal distribution. Equation (1) assumes that all individuals are drawn from the same population. The means of the latent growth factors α show the average development of the measurement y_i across p panel waves within a homogeneous population.

2.2 Growth mixture models

Growth mixture models can relax the assumption of a homogeneous population and can give information about parameter differences across unobserved subpopulations.

Instead of considering individual variation of single means of the vector η the growth mixture model allows different classes of individuals to vary around different means. Classes are introduced by a latent categorical variable where the categories represent the unobserved heterogeneity of the data (Muthén and Shedden 1999):

$$y_{ik} = \Lambda_k \eta_{ik} + \epsilon_{ik} \quad (6)$$

The growth mixture model (abbreviated GMM) in (6) allows the estimation of $k = 1, \dots, K$ latent classes. The probability density function for the GMM is a finite mixture of normal distributions:

$$f(y_i) = \sum_{k=1}^K \pi_k \phi_k[y_i; \mu_k(\theta_k) \Sigma(\theta_k)] \quad (7)$$

π_k is the unconditional probability that a measurement belongs to latent class k , ϕ_k is the multivariate probability density function for latent class k . $\mu_k(\theta_k)$ represents the model-implied mean vector given by

$$\mu_k(\theta_k) = \Lambda_k \alpha_k \quad (8)$$

and $\Sigma_k(\theta_k)$ is the model-implied covariance matrix given by

$$\Sigma_k(\theta_k) = \Lambda_k \Psi_k \Lambda'_k + \Theta_k \quad (9)$$

In an unconditional mixture model the latent variables η are only described by their class specific means α_k and variances Ψ_k . A conditional mixture model includes exogenous latent variables ξ_n representing the observed heterogeneity of the data. The relation between ξ_n and the categorical class variable c is given by a multinomial logistic regression equation:

$$\text{logit}(\pi_k) = \alpha_k + \Gamma_k \xi_n \quad (10)$$

with $\pi_k = P(c_k = k | \xi_n)$. Γ_k is a $(K - 1) \times q$ -parameter matrix containing regression coefficients of K classes on ξ_n .

The simplest mixture model is *latent class growth analysis* (abbreviated LCGA), which is a submodel of (6) assuming that growth factors η are fixed instead of random effects, i.e. they have zero variances within the particular classes ($\Psi = 0$). So, classes are treated as homogeneous with respect to their development. Two major advantages are emphasized (Muthén 2004, p. 350): It can be used to find cutpoints in the within-class variation on the growth factors and it can serve as a starting point for the more general GMM.

Growth mixture models are estimated by maximizing the log-likelihood function within the admissible range of parameter values given classes and data. *Mplus* employs the EM-algorithm for maximization (Dempster et al. 1977; Muthén and Shedden 1999). For a given solution, each individual's probability of membership in each class is estimated. Individuals can be assigned to the classes by calculating the posterior probability that an individual i belongs to a given class k . Each individual's

posterior probability estimate for each class is computed as a function of the parameter estimates and the values of the observed data.

Standard errors of estimates are asymptotically correct if the underlying mixture model is the true model. In general, test statistics require well-defined classes in a mixture model. In mixture models a k class model is not nested within a $k + 1$ class model. Therefore, χ^2 -differences cannot be used for statistical tests. Usually, the Bayesian Information Criterion (BIC; Schwarz 1978) are used for model comparisons which was found to perform best for growth mixture models in a simulation study by Nylund et al. (2007). The model with the smallest BIC is accepted within model comparisons.

2.3 Special cases for count data

2.3.1 Poisson and zero-inflated Poisson model

When the response variable under study is a count, the Poisson regression as a special case of the generalized linear model are often applied. Let y_i be the number of observed count occurrences, x_i be the vector of covariates and v_i be the expected number of counts.¹ The number of events in an interval of a given length is Poisson distributed and the Poisson regression model can be formulated via a log link function (Hilbe 2007, p. 39; Greene 2008, p. 585):

$$\Pr(y_i|x_i) = \exp(-v_i)v_i^{y_i}/y_i! \quad (11)$$

with $v_i = \exp(\alpha + x_i'\beta)$. β is the vector of regression coefficients. The conditional mean function of the Poisson distribution is $E(y_i|x_i) = v_i$ with its equidispersion $\text{Var}(y_i|x_i) = v_i$.

If the number of zeros in the count variable are very large, a variant of the Poisson regression model is more appropriate: the so-called *zero-inflated Poisson model* (abbreviated ZIP). The ZIP model combines the Poisson regression model in (11) with a logit model to cover the zero inflation in the count variable (Lambert 1992):

$$\Pr(y_i|x_i) = \begin{cases} \pi_i + (1 - \pi_i)\exp(-v_i) & \text{for } y_i = 0 \\ (1 - \pi_i)\frac{\exp(-v_i)v_i^{y_i}}{y_i} & \text{for } y_i \geq 1 \end{cases} \quad (12)$$

π is the probability of being an extra zero. A growth mixture model with two parts are estimated simultaneously when zero inflation of the data is assumed: the first part contains the Poisson model of the measurements with values of zero and above and the second part refers to the logit model of the measurements with values of zeros across the panel waves. The Poisson and the ZIP model can be applied with the program Mplus (starting with version 3).

¹Note that usually λ is used in the Poisson model instead of v . But here the authors are using λ for parameters (factor loadings) congruent to the terminology in structural equation models.

2.3.2 Negative binomial and zero-inflated negative binomial model

If the assumption of equidispersed data does not hold, the negative binomial regression model can be employed by introduction of latent heterogeneity in the conditional mean of the Poisson model (Hilbe 2007, p. 207; Greene 2008, p. 586):

$$\Pr(y_i|x_i, \epsilon_i) = \exp(\alpha + x'_i\beta + \epsilon_i) = h_i v_i \quad (13)$$

where $h_i = \exp(\epsilon_i)$ is assumed to have a one parameter gamma distribution, $G(\theta, \theta)$ with mean equal to 1 and variance $\kappa = 1/\theta$. The negative binomial distribution can be obtained by integrating h_i out of the joint distribution. The conditional mean function is still $E(y_i|x_i) = v_i$ while overdispersion can be obtained from the latent heterogeneity with the variance function $\text{Var}(y_i|x_i) = v_i^2[1 + (1/\theta)]$. Because of the quadratic term for v_i , the negative binomial model was labeled NB-2. Other variance functions lead to other types of negative binomial models (Hilbe 2007, p. 78).

With large number of zeros in the count variable, the so-called *zero-inflated negative binomial model* (abbreviated ZINB) is more appropriate. Similar to the ZIP model the ZINB model combines the negative binomial regression model with a logit model to cover the zero inflation in the count variable (Hilbe 2007, p. 160f.).²

3 Study, sample and variables

The application of different growth curve and growth mixture models was conducted with data from the ongoing panel study *Crime in the Modern City* (*CrinoC*).³ Eight annual panel waves had been collected between 2002 and 2009 from a sample of adolescents with a mean age of 13 (7th grade) in the initial survey ($n = 3411$). The sample was drawn from schools in Duisburg, an industrial city of approximately 500000 inhabitants located in the western part of the Ruhr area. Self-administered questionnaires were completed in school classes. After leaving the school at the end of the 10th grade, adolescents had to be contacted by mail or personally at home.

Data for the following analyses stem from a five-wave panel data set covering the period from late childhood to late adolescence (age 13–17). Included are 1552 adolescents who participated in all panel waves and comprised 45.5% of the initial sample. This indicates a common but nevertheless considerable attrition that can partly be ascribed to the method used for panel construction (Pöge 2005)⁴ and to certain characteristics of the respondents. As can be seen from Table 1, the distributions of the

²Note that several software implementations allow the zero-inflated binary process to be either probit or logit (for details, Hilbe 2007, p. 174). The negative binomial (NB-2) and the ZINB model can be applied with the program *Mplus* (starting with version 5.2).

³Detailed information about the study can be obtained from the webpage www.crimoc.org.

⁴To avoid high refusal rates in the initial sample by collecting respondents' names and addresses, respondents were asked to fill out and reproduce an individual and unique code year by year. The code was generated by questions regarding particular individual and rememberable characteristics (e.g., first letter of mothers name). However, participants to a large degree seem to differ in their ability to reproduce the code correctly.

Table 1 Panel (2002–2006) and initial cross sectional (2002) sample

| | Panel | | Cross section | |
|-------------------|-------|---------|---------------|---------|
| | n | (%) | n | (%) |
| ♂ | 642 | (41.4) | 1728 | (50.7) |
| ♀ | 910 | (58.6) | 1679 | (42.2) |
| School (high) | 392 | (25.3) | 778 | (22.8) |
| School (medium 2) | 517 | (33.3) | 1064 | (31.2) |
| School (medium 1) | 373 | (24.0) | 806 | (23.6) |
| School (low) | 270 | (17.4) | 763 | (22.4) |
| Total | 1552 | (100.0) | 3411 | (100.0) |

Note: Educational level of schools given in brackets

Table 2 Descriptives for annual self-reported delinquency

| Age | | \bar{y} | s^2 | s^3 | s^4 | %zero |
|-----|-------|-----------|---------|--------|---------|-------|
| 13 | t_1 | 2.688 | 139.393 | 9.259 | 114.448 | 74.98 |
| 14 | t_2 | 6.264 | 615.899 | 8.007 | 99.100 | 68.90 |
| 15 | t_3 | 5.202 | 593.146 | 12.446 | 235.413 | 71.46 |
| 16 | t_4 | 4.216 | 451.517 | 9.908 | 131.694 | 76.34 |
| 17 | t_5 | 4.105 | 540.799 | 9.281 | 101.293 | 82.29 |

\bar{y} = mean; s^2 = variance; s^3 = skewness; s^4 = kurtosis

panel respondents by gender and educational level are somewhat biased compared to the distributions in the cross-sectional sample from 2002. This indicates that girls as well as high educated respondents are overrepresented in the panel data used for analyses.

Several applications of growth mixture models with data from this investigation and a previous pilot study have been published Reinecke (2006a, 2006b, 2007, 2008). While these analyses were focused on the annual summed prevalence rates, i.e. the versatility of adolescents delinquent behavior, the current analyses are focused on the incidence rates to directly study the development of the frequency, i.e. the intensity of delinquent behavior. The annual incidence rate is a composite measure of self-reported delinquency. Respondents were asked to give the number of delinquent behaviors committed during the last 12 months for 16 different offenses separately. The offenses were theft of and out of cars, theft out of a vending machine, theft of bicycles, other thefts, burglary, shoplifting, fencing, robbery, purse snatching, assault with and assault without a weapon, graffiti, scratching, damage to property and drug dealing. The overall annual incidence rate is given as the sum of the rates for the 16 behaviors. Table 2 contains descriptive information about the annual composite measures of self-reported delinquency. After a peak at age 14 (t_2) the mean frequency of delinquent activity is constantly declining. The distributions are characterized by rel-

atively high proportions of persons who do not report any delinquent activity (% zeros) and relatively few persons with extreme values. Large variances, skewness and kurtosis are further characteristics which are typical for behavioral data relying on rare events. Thus, the measures of annual self-reported delinquency can be treated as overdispersed count data with an inflation of zeros.

The following analyses are divided into three parts: first, techniques of latent growth curve models will be used to specify the observed outcome as a function of time (respective age) alone and to check for potential variations around the growth factors means. Second, latent class growth and growth mixture model specifications will be applied to the data. Furthermore, the best fitting solutions two alternative modeling approaches (zero-inflated Poisson and zero-inflated negative binomial) will be compared with regard to the differences and similarities of assigning individuals to latent classes. Third, the best growth mixture model will be enlarged by adding covariates in order to give auxiliary information for a more precise classification and to incorporate potential predictors of the particular latent class distributions.

4 Latent growth models

The basic functional form of the growth process for all subsequent (unconditional) latent growth and growth mixture models is given in (14), (15), and (16):⁵

$$\begin{bmatrix} y_{1ik} \\ y_{2ik} \\ y_{3ik} \\ y_{4ik} \\ y_{5ik} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \\ 1 & 4 & 16 \end{bmatrix} \begin{bmatrix} \eta_{1ik} \\ \eta_{2ik} \\ \eta_{3ik} \end{bmatrix} + \begin{bmatrix} \epsilon_{1ik} \\ \epsilon_{2ik} \\ \epsilon_{3ik} \\ \epsilon_{4ik} \\ \epsilon_{5ik} \end{bmatrix} \quad (14)$$

According to the development of the mean level of self-reported delinquency (Table 2) the specification of a quadratic term was deemed necessary to account for non-linear change. The growth factor loadings were accordingly fixed in matrix Λ_k . The latent growth factors are described as a function of their means in vector α_k and residual parameters in vector ζ_k :

$$\begin{bmatrix} \eta_{1ik} \\ \eta_{2ik} \\ \eta_{3ik} \end{bmatrix} = \begin{bmatrix} \alpha_{1k} \\ \alpha_{2k} \\ \alpha_{3k} \end{bmatrix} + \begin{bmatrix} \zeta_{1ik} \\ \zeta_{2ik} \\ \zeta_{3ik} \end{bmatrix} \quad (15)$$

Matrix Ψ_k contains the variances and covariances of the latent growth factors:

$$\Psi_k = \begin{bmatrix} \psi_{11k} & & \\ \psi_{21k} & \psi_{22k} & \\ \psi_{31k} & \psi_{32k} & \psi_{33k} \end{bmatrix} \quad (16)$$

⁵ $K = 1$ for latent growth models.

Table 3 Comparison of different latent growth model specifications

| Model | Random effects | Parameters | Log-likelihood | BIC |
|-------|----------------|------------|----------------|-----------|
| ZIP1 | – | 6 | −39092.919 | 78229.921 |
| ZIP2 | <i>I</i> | 7 | −23596.340 | 47244.112 |
| ZIP3 | <i>IS</i> | 9 | −17714.575 | 35495.276 |
| ZIP4 | <i>ISQ</i> | 12 | −16007.225 | 32102.617 |
| ZINB1 | – | 11 | −10924.769 | 21930.358 |
| ZINB2 | <i>I</i> | 12 | −10206.846 | 20501.860 |
| ZINB3 | <i>IS</i> | 14 | −10164.167 | 20431.195 |
| ZINB4 | <i>ISQ</i> | 17 | −10156.666 | 20438.237 |

I = intercept; *S* = linear slope; *Q* = quadratic slope

Table 4 Estimated random effects for ZINB models

| Parameter | ZINB2 | | ZINB3 | | ZINB4 | |
|-----------|-------|-----------|-------|-----------|-------|-----------|
| | Est. | (z-value) | Est. | (z-value) | Est. | (z-value) |
| ψ_I | 8.522 | (18.645) | 9.423 | (12.434) | 6.687 | (10.009) |
| ψ_S | – | – | 0.551 | (7.044) | 2.491 | (5.496) |
| ψ_Q | – | – | – | – | 0.160 | (4.796) |

Model specifications varied from fixing the variances and covariances of the latent growth factors to zero (fixed effects model) to a completely random effect specification (random effects model).⁶ Due to the large amount of non-delinquent adolescents the models were specified as zero-inflated. The possibility of significant overdispersion to the outcomes variables was considered by testing the zero-inflated Poisson (ZIP) against the zero-inflated negative binomial model (ZINB).

Results clearly show that, according to the log-likelihood and the Bayesian Information Criterion (BIC), the ZINB models outperform the ZIP models (Table 3). The additional consideration of overdispersion seems more suitable to represent the outcome variable as negative binomial distributed. Correspondingly, the estimated dispersion parameters for the outcomes in all ZINB models have values >0 ($p < 0.05$). Furthermore, within the ZINB models the fixed effects specification (ZINB1) is outperformed by the random effects models (ZINB2 to ZINB4). According to the BIC, the model with random effects for the intercept and linear slope and fixed effect for the quadratic slope performs best (ZINB3). Parameter estimates for the growth factor variances are significant in all ZINB model specifications with random effects (Table 4). Even the small variance of the quadratic slope in the model ZINB4 is signifi-

⁶In the fixed effects model the variances of the intercept, the linear and quadratic slope were fixed to zero, assuming that all individuals follow a unique developmental trajectory. In the random effects model the variances of the growth factors were estimated, assuming that all individuals follow a unique developmental trajectory, but may vary in their initial levels and the extent of change indicated by the linear and the quadratic slope.

Table 5 Comparison of latent class growth models (ZINB)

| Model | Classes | Parameters | Log-likelihood | BIC |
|--------------------|----------|------------|-------------------|------------------|
| LCGA (ZINB) | 2 | 15 | -10326.262 | 20762.734 |
| LCGA (ZINB) | 3 | 19 | -10169.237 | 20478.073 |
| LCGA (ZINB) | 4 | 23 | -10142.068 | 20453.123 |
| LCGA (ZINB) | 5 | 27 | -10116.018 | 20430.414 |
| LCGA (ZINB) | 6 | 31 | -10098.193 | 20424.152 |
| LCGA (ZINB) | 7 | 35 | -10087.260 | 20431.676 |
| LCGA (ZINB) | 8 | 39 | -10076.222 | 20438.989 |

cant although the BIC is slightly higher than the BIC in the previous model ZINB3. Hence, the results indicate that individuals randomly vary around one developmental trajectory that applies to all individuals in the sample.

5 Latent class growth and growth mixture models

The significant variation around the growth factor means from the latent growth analysis might be an indication of possible unobserved heterogeneity in adolescents' development of delinquency. The hypothesis to detect more than one distinct group of individuals was analyzed by applying two different parameterizations of mixture models to the data. First, with a latent class growth analysis (LCGA) models with increasing numbers of classes representing distinctive developmental trajectories were tested. Note that LCGA allows no variations of the growth factors. Furthermore, the models were estimated with respect to the zero-inflated negative binomial distribution of the outcome variables. Log-likelihood and the BIC values were used to identify the sufficient number of classes.

Table 5 gives an overview of the model results. The resulting best LCGA model has six classes with the lowest BIC value (20424.152). The model estimated dispersion parameters for the five outcomes (panel waves) in the six-class model are $t_1 = 2.220$ ($z = 1.862$), $t_2 = 2.337$ ($z = 6.338$), $t_3 = 2.594$ ($z = 6.343$), $t_4 = 3.403$ ($z = 10.092$) and $t_5 = 6.621$ ($z = 6.613$).⁷ The estimated degree of overdispersion again supports the estimation of the model on the basis of the negative binomial model assumptions.

According to the estimated model the largest class represents a group of adolescents who are nearly not involved in any delinquent activity during the observed period (non offenders, 43.7%). The second largest class are individuals who are supposed to commit very few delinquent acts (low level offenders, 18.7%). The third largest class is starting from a low level of delinquency, followed by a slight increase until a peak at age 15. A likewise slight decrease can be observed from there on until the initial level is attained again (adolescence limited offenders, 15.8%). The

⁷Dispersion parameters were held equal across classes.

Table 6 Estimated means and residuals for the six-class LCGA model

| Class | | t_1 | t_2 | t_3 | t_4 | t_5 |
|-----------------------|------------|--------|--------|--------|--------|--------|
| Non offenders | Est. means | 0.025 | 0.009 | 0.006 | 0.008 | 0.022 |
| | Residuals | 0.000 | 0.001 | 0.000 | 0.000 | 0.000 |
| Low level | Est. means | 0.751 | 0.758 | 0.564 | 0.380 | 0.231 |
| | Residuals | -0.033 | 0.063 | -0.008 | -0.042 | 0.024 |
| Adolescence limited | Est. means | 2.160 | 5.490 | 7.002 | 5.494 | 2.652 |
| | Residuals | -0.012 | -0.157 | 0.730 | -0.780 | 0.249 |
| High level/persistent | Est. means | 19.287 | 37.139 | 42.860 | 36.344 | 22.645 |
| | Residuals | -2.123 | 9.269 | -5.701 | -4.751 | 4.135 |
| Late starters | Est. means | 0.027 | 0.123 | 0.523 | 2.497 | 13.380 |
| | Residuals | -0.002 | -0.004 | 0.055 | -0.263 | 0.894 |
| Early decliners | Est. means | 8.123 | 9.209 | 0.856 | 0.008 | 0.000 |
| | Residuals | -0.097 | 0.006 | -0.001 | 0.000 | 0.000 |

fourth largest class is characterized by individuals with a high and partial persistent level of delinquent activity (high level and persistent offenders, 10.2%). The two smallest classes are of special interest for the prediction of chronic or intensive delinquent behavior over the life-course. Both of them challenge the assumption that the early onset of delinquent behavior is one of the best characteristics for the prediction of long-term criminal careers (Farrington et al. 1990; Odgers et al. 2007): the second smallest class represents a pathway of individuals with nearly no estimated delinquent activity for the first three years of the observation, but then engage in delinquency arriving at a comparatively high level at the end of adolescence (late starters, 6.0%). Reversely, the smallest group shows a remarkable level of delinquency at age 13, then it rapidly declines to a near zero level (early decliners, 5.6%). Both trajectories would not be expectable from early onset prediction. Table 6 displays the model estimated means and residuals, indicating the developmental trajectories for the six classes based on the estimated posterior probabilities.⁸ Especially for the class of high level offenders the residuals indicate considerable degrees of over- and underestimation of the means. The other residuals indicate at most small differences between estimated and observed means.

Based on the results of the LCGA, a growth mixture model with random effects was specified and tested. Previous analyses have shown that variances of the linear

⁸For further analyses it may be desirable to use a classification variable that assigns every individual to exactly one class by the individuals highest probability of class membership. This absolute and determined classification may differ from the solution based on estimated posterior probabilities, especially when the separation of the classes lacks accuracy. A criterion to assess the quality of the classification is the entropy criterion $E(k)$ (Celeux and Soromenho 1996). For example, the entropy for the six-class LCGA model is $E(k) = 0.661$. However, despite the indistinctness in classification such solutions can be suitable for substantive interpretation.

Table 7 Comparison of growth mixture models (ZINB)

| Model | Classes | Parameters | Log-likelihood | BIC |
|-------------------|----------|------------|-------------------|------------------|
| GMM (ZINB) | 2 | 16 | -10158.691 | 20434.938 |
| GMM (ZINB) | 3 | 20 | -10121.995 | 20390.936 |
| GMM (ZINB) | 4 | 24 | -10093.843 | 20364.022 |
| GMM (ZINB) | 5 | 28 | -10083.137 | 20371.998 |
| GMM (ZINB) | 6 | 32 | -10074.556 | 20384.226 |
| GMM (ZINB) | 7 | 36 | -10070.231 | 20404.965 |

and quadratic growth factor could be fixed to zero while the intercept variance was estimated but set equal across classes (Mariotti and Reinecke 2010). Table 7 shows the results with up to seven classes. Like the LCGA models, the GMM models were estimated with respect to the zero-inflated negative binomial distribution. According to the smallest BIC value the best model has four classes. Models with five classes and above show considerable higher BIC values. The log-likelihood differences diminish with an increase of the number of classes. Just as for the LCGA model with six classes the model estimated dispersion parameters indicate significant levels ($t_1 = 3.250$ ($z = 3.385$), $t_2 = 2.253$ ($z = 9.710$), $t_3 = 2.558$ ($z = 10.334$), $t_4 = 2.668$ ($z = 7.998$) and $t_5 = 5.244$ ($z = 7.406$)). The estimated intercept variance turned out to be significant ($\psi_I = 2.076$ ($z = 4.732$)) indicating remarkable variation around the base level of self-reported delinquent behavior.

Again, the largest class represents individuals with nearly no delinquent activity (non offenders, 43.1%). The second largest class is estimated to have a mean trajectory of persistent and frequent delinquent activity (high level and persistent offenders, 28.6%). The third largest class starts with a quite remarkable level of delinquent activity at age 13 and even enhances the estimated mean frequency at age 14. Afterwards, delinquency constantly decreases to lower levels throughout middle adolescence (adolescence limited, 18.0%). The smallest class represents the developmental path of individuals who engage in delinquent activity by the end of adolescence (late starters, 8.7%).⁹ Table 8 displays the model estimated means and residuals for the four classes based on the estimated posterior probabilities. Here, the estimated mean values for the class of high level offenders seem to suffer from substantive underestimation. Additionally, the estimated mean for the t_5 in the late starters class is somewhat underestimated. Despite these notable discrepancies between estimated and observed means the residuals are quite low.

A comparison of the fit and information criteria of the LCGA model with six classes and the GMM model with four classes shows lower log-likelihood (-10098.193 vs. -10093.843) and BIC values (20424.152 vs. 20364.022) for the GMM model. Especially the difference in the log-likelihood values is small. Based

⁹ Although the model with four classes is statistically the favorable solution, the five-class model can be of at least the same interest for substantive research questions. The fifth class represents an early withdraw from delinquent activity (so-called early deliners). The decision about the correct number of classes on the basis of the BIC is a statistical one and should be reflected with substantive arguments. Given the methodological character of this paper, the four-class solution is discussed here.

Table 8 Estimated means and residuals for the four-class GMM model

| Class | | t_1 | t_2 | t_3 | t_4 | t_5 |
|---------------------------|------------|--------|--------|--------|--------|--------|
| Non offenders | Est. means | 0.101 | 0.019 | 0.009 | 0.012 | 0.042 |
| | Residuals | 0.008 | -0.001 | 0.000 | 0.000 | 0.000 |
| High level/ persistent | Est. means | 8.545 | 16.733 | 21.087 | 18.082 | 10.551 |
| | Residuals | -2.618 | -0.512 | -5.108 | -4.412 | 0.277 |
| Adolescence limited | Est. means | 5.770 | 8.609 | 2.835 | 0.218 | 0.004 |
| | Residuals | -0.962 | -0.432 | 0.060 | -0.016 | 0.001 |
| Late starters | Est. means | 0.040 | 0.143 | 0.566 | 2.664 | 14.861 |
| | Residuals | 0.001 | -0.026 | 0.079 | -0.319 | -3.487 |

Table 9 Comparison of LCGA and GMM solutions (ZINB)

| | | LCGA | | | | | | |
|-------|------|------|-----|-----|-----|-----|-----|-------|
| | | (1) | (2) | (3) | (4) | (5) | (6) | Total |
| GMM | (7) | 772 | | | | | 12 | 784 |
| | (8) | | 117 | 190 | 10 | | 66 | 383 |
| | (9) | | 16 | 32 | | 68 | 175 | 291 |
| | (10) | | | | 57 | | 37 | 94 |
| Total | | 772 | 133 | 222 | 67 | 68 | 290 | 1552 |

Note: (1) non offenders, (2) persistent offenders, (3) adolescence limited, (4) late starters, (5) early decliners, (6) low level offenders, (7) non offenders, (8) persistent offenders, (9) adolescence limited, (10) late starters

on the most likely class membership of individuals the four-class GMM and six-class LCGA solutions can be crosstabulated in order to compare for differences in the classification (Table 9). The non offending individuals have been almost equally classified within both modeling approaches. Moreover, the GMM adolescence limited class consists mostly of individuals from the LCGA low level and early decline classes. The GMM late starters class contains most of the LCGA late starters. Finally, the GMM high level and persistent class has nearly tripled its size mainly by absorbing 256 individuals from the LCGA adolescence limited and low level classes. Altogether, the six distinct classes—as a result of the fixed effects model (LCGA)—are now represented by fewer and more general classes in the GMM. This decrease of the number of classes in the particular mixture model specification was expected and is in accordance with other applications comparing LCGA and GMM models (Muthén 2004; Kreuter and Muthén 2008a).

By inspecting the observed individual curves for the three offending classes of the four-class GMM model it is obvious that because of the massive absorption of the individuals from different classes of the LCGA model, the high level and persistent group is the most heterogeneous of the four classes (Fig. 2). The displayed subset of 72 randomly selected curves of individuals who are classified as high level and




Fig. 2 Observed individual trajectories for the three offender classes (four-class GMM model)

Table 10 Comparison of unconditional and conditional four-class GMM model (ZINB)

| | | Unconditional | | | | |
|-------------|-----|---------------|-----|-----|-----|-------|
| | | (1) | (2) | (3) | (4) | Total |
| Conditional | (1) | 775 | 3 | | 13 | 791 |
| | (2) | 2 | 327 | 3 | 1 | 333 |
| | (3) | 7 | 50 | 288 | | 345 |
| | (4) | | 3 | | 80 | 83 |
| Total | | 784 | 383 | 291 | 94 | 1552 |

Note: (1) non offenders, (2) persistent offenders, (3) adolescence limited, (4) late starters

persistent, is characterized by numerous differing “zick–zack” patterns of drifting in and out of delinquent activity during the observational period. Although these heterogeneous pathways for the most part display high levels of delinquency at a particular point, a more or less consistent pattern of *persistent* high level offending is hardly observable. The observed curves of the individuals classified as adolescence limited or late starters to a greater extent correspond with the estimated mean trajectories of the classes. Also, within the adolescence limited class several trajectories of the absorbed early decliners from the LCGA model can be identified.

In a further step the four-class GMM model was again estimated with additional covariates in order to check if the four-class solution remains stable. This enhanced model specification uses multinomial logistic regression to predict class membership from additional information of covariates (cf. (10)) and to further exhibit the profile of the individuals in the classes (Muthén 2002). For this analysis additional information on gender and educational level (as displayed in Table 1) were used. The resulting four-class conditional GMM has a log-likelihood value of -10047.658 and a BIC of 20315.734. Both values are considerably lower than for the unconditional model (-10093.843 and 20364.022). The model estimated variance in the intercept factor again is significant ($\psi_1 = 1.929$ ($z = 6.865$)) as well as the estimated dispersion parameters ($t_1 = 3.195$ ($z = 3.420$)), $t_2 = 2.300$ ($z = 9.648$)), $t_3 = 2.689$ ($z = 10.450$)), $t_4 = 2.721$ ($z = 7.592$) and $t_5 = 4.673$ ($z = 7.619$)).

The classification of individuals based on posterior probabilities resulted in equally shaped distinct trajectories with similar class proportions compared to the unconditional four-class model (non offenders, 45.0%; high level and persistent offenders, 25%; adolescence limited, 22.5%; late starters, 7.5%). The classification results of the unconditional and conditional model are compared in Table 10 based on the most likely class membership of the respondents. Three aspects are noteworthy: First, in the conditional GMM model 50 persons were classified into the adolescence limited class which have to be in the high level and persistent class of the unconditional GMM model. Second, 13 persons switched from the late starter class in the unconditional model to the class of non offenders in the conditional model. Third, the overall stability of the classification is indicated by high values in the diagonal of the table. All in all, changes in class sizes are moderate.

Table 11 displays the results of the multinomial logistic regression of class membership on the categories of gender and educational level. For boys on a low edu-

Table 11 Odds (e^{logit}) and class probabilities for the conditional four-class GMM model (ZINB)

| | School (low) ♂ | | School (med 1) ♂ | | School (med 2) ♂ | | School (high) ♂ | |
|---------------|--------------------|-------|--------------------|-------|--------------------|-------|--------------------|-------|
| | e^{logit} | Prob. | e^{logit} | Prob. | e^{logit} | Prob. | e^{logit} | Prob. |
| Non offenders | 1.000 | 0.256 | 1.000 | 0.295 | 1.000 | 0.338 | 1.000 | 0.382 |
| High level | 1.707 | 0.436 | 1.363 | 0.402 | 1.089 | 0.368 | 0.868 | 0.333 |
| Ado. lim. | 0.818 | 0.209 | 0.675 | 0.199 | 0.546 | 0.185 | 0.451 | 0.173 |
| Late starters | 0.387 | 0.099 | 0.353 | 0.104 | 0.322 | 0.109 | 0.293 | 0.112 |

| | School (low) ♀ | | School (med 1) ♀ | | School (med 2) ♀ | | School (high) ♀ | |
|---------------|--------------------|-------|--------------------|-------|--------------------|-------|--------------------|-------|
| | e^{logit} | Prob. | e^{logit} | Prob. | e^{logit} | Prob. | e^{logit} | Prob. |
| Non offenders | 1.000 | 0.460 | 1.000 | 0.509 | 1.000 | 0.556 | 1.000 | 0.602 |
| High level | 0.424 | 0.195 | 0.338 | 0.172 | 0.270 | 0.150 | 0.216 | 0.130 |
| Ado. lim. | 0.635 | 0.292 | 0.524 | 0.266 | 0.433 | 0.241 | 0.357 | 0.216 |
| Late starters | 0.115 | 0.053 | 0.105 | 0.053 | 0.096 | 0.053 | 0.087 | 0.052 |

cational level the probability to be classified into one of the three offending classes is about 0.75 with the highest single probability for the high level class (0.44). But males attending school types with the highest educational level have a probability of about 0.62 to be classified into one of the three offending classes. The probability for the adolescence limited class decreases with the educational level of the schools. Interestingly, even though to a small extent, the probability for the late starter group of boys increases with higher educational levels.

Girls who visit schools with high educational level are most likely to be classified as non offenders (0.60). Even for girls in schools with the lowest educational levels the highest class probabilities appear for the class of non offenders (0.46). The highest probability to be in one of the offending classes emerges for the adolescence limited class, but decreases with higher levels of education. The low probability of the high level class also decreases. The probability of the late starters class of girls remains stable across school types. Altogether, the main differences seem to exist between boys and girls, pointing to gender as a relatively strong predictor of class membership.

6 Discussion

The general framework of growth mixture modeling outlined by Muthén (2002, 2004, 2008) integrates continuous and categorical approaches of longitudinal data analysis. A growth mixture model contains a growth curve model formalized with structural equations. A categorical variable covers the mixture distribution via a latent class model. Furthermore, a multinomial regression model formalizes the relationships between exogenous time-invariant variables and the latent class variable. A popular submodel, explored and discussed by Nagin and Land (1993) and Nagin (1999) is LCGA, in which the variances of the growth curve parameters are fixed to zero. Due to an easier estimation of the parameters, LCGA is computationally less demanding

and thus useful for a first evaluation of the unobserved heterogeneity of the data. If count data with large amounts of zeros are analyzed, the outcome variables can be assumed as zero-inflated Poisson or zero-inflated negative binomial distributed. The latter one considers highly overdispersed distributions.

Data from a five-wave panel study of adolescents have been used to study unobserved heterogeneity in the development of deviant and delinquent behavior. In a first step the outcome variable has been analyzed by means of a latent quadratic growth model under zero-inflated Poisson and zero-inflated negative binomial specifications. Due to the overdispersed data, model variants with the negative binomial specification have always better model fits compared to those ones assuming Poisson distributions.

With latent class growth analysis (LCGA) models up to eight classes have been tested. According to the BIC, the model with six classes performed best. All classes can be interpreted substantially. But LCGA treats the growth curve variables as fixed effects and thus does not account for possible variation within the classes. Possible overlaps of individual trajectories from different classes are therefore ignored.

Within GMM the variance of the intercept was estimated for models up to seven classes. According to the BIC, the model with four classes performed best. Similar to analyses of Muthén (2004) and Kreuter and Muthén (2008a) GMM results in less classes than LCGA. The consideration of possible overlaps in trajectories leads to a better substantive interpretation of the development of delinquency. Additional variations of the linear slope and the quadratic slope did not result in further model improvements albeit model estimations problems increased with additional parameters to be estimated.

Decisions about the correct number of classes in growth mixture models can also involve a likelihood ratio-based method for testing $k - 1$ classes against k classes developed by Lo et al. (2001) (LMR-LRT) or the bootstrapped likelihood ratio test (BLRT) proposed by McLachlan and Peel (2000). The performance of these tests for non normal outcomes is unclear (Jeffries 2003) and the implications for applications has not been discussed yet.

A next step in analysis could be to test a non-parametric specification of the growth mixture model (Kreuter and Muthén 2008a, 2008b; Muthén and Asparouhov 2008). While the basic GMM assumes a specific (normal) distribution for the random effects, the non-parametric version does not make any such assumption. However, a first inspection of the individual intercept factor scores for the three offender classes from the four-class GMM model presented in this paper provided no clear evidence for a non-normally distributed variation around the estimated intercept.

References

- Boers, K., Reinecke, J., Seddig, D., Mariotti, L.: Explaining the development of adolescent violent delinquency. *Eur. J. Criminol.* **7**, 499–520 (2010)
- Celeux, G., Soromenho, G.: An entropy criterion for assessing the number of clusters in a mixture model. *J. Classif.* **13**, 195–212 (1996)
- Curran, P.J., Muthén, B.: The application of latent curve analysis to testing developmental theories in intervention research. *Am. J. Community Psychol.* **27**, 567–595 (1999)
- Dempster, A.P., Laird, N.M., Rubin, D.B.: Maximum likelihood from incomplete data via the EM algorithm. *J. R. Stat. Soc. B* **39**(1), 1–38 (1977)

- Duncan, T.E., Duncan, S.C., Strycker, L.A.: An Introduction to Latent Variable Growth Curve Modeling. Concepts, Issues and Applications, 2nd edn. Lawrence Erlbaum, Mahwah (2006)
- Farrington, D.P., Loeber, R., Elliott, D.S.: Advancing knowledge about the onset of delinquency and crime. In: Lahey, B., Kazdin, A. (eds.) Advances in Clinical Child Psychology, vol. 13, pp. 283–342. Plenum Press, New York (1990)
- Greene, W.: Functional forms for the negative binomial model for count data. *Econ. Lett.* **99**, 585–590 (2008)
- Hilbe, J.M.: Negative binomial regression. Cambridge University Press, Cambridge (2007)
- Jeffries, N.O.: A note on testing the number of components in a normal mixture. *Biometrika* **90**, 991–994 (2003)
- Kreuter, F., Muthén, B.: Analyzing criminal trajectory profiles: bridging multilevel and group-based approaches using growth mixture modeling. *J. Quant. Criminol.* **24**, 1–31 (2008a)
- Kreuter, F., Muthén, B.: Longitudinal modeling of population heterogeneity: methodological challenges to the analysis of empirically derived criminal trajectory profiles. In: Hancock, G.R., Samuelsen, K.M. (eds.) Advances in Latent Variable Mixture, Modeling, pp. 53–75. Information Age Publishing, Charlotte (2008b)
- Lambert, D.: Zero-inflated Poisson regression with an application to defects in manufacturing. *Technometrics* **34**, 1–13 (1992)
- Lo, Y., Mendell, N.R., Rubin, D.B.: Testing the number of components in a normal mixture. *Biometrika* **88**(3), 767–778 (2001)
- Mariotti, L., Reinecke, J.: Wachstums- und Mischverteilungsmodelle unter Berücksichtigung unbeobachteter Heterogenität: Empirische Analysen zum delinquenten Verhalten Jugendlicher in Duisburg. Institut für sozialwissenschaftliche Forschung e.V., Münster (2010)
- McArdle, J.J.: Dynamic but structural equation modeling of repeated measures data. In: Nesselroade, J.R., Cattell, R.B. (eds.) Handbook of Multivariate Experimental Psychology, 2nd edn., pp. 561–614. Plenum, New York (1988)
- McArdle, J.J., Epstein, D.: Latent growth curves within developmental structural equation models. *Child Dev.* **58**, 110–133 (1987)
- McLachlan, G., Peel, D.: Finite Mixture Models. Wiley, New York (2000)
- Meredith, M., Tisak, J.: Latent curve analysis. *Psychometrika* **55**(1), 107–122 (1990)
- Muthén, B.: Analysis of longitudinal data using latent variable models with varying parameters. In: Collins, L.M., Horn, J.L. (eds.) Best Methods for the Analysis of Change: Recent Advances, Unanswered Questions, Future Directions, pp. 1–17. APA, Washington (1991)
- Muthén, B.: Latent variable modeling with longitudinal and multilevel data. *Sociol. Method.* **27**, 453–480 (1997)
- Muthén, B.: Latent variable mixture modeling. In: Marcoulides, G.A., Schumacker, R.E. (eds.) New Developments and Techniques in Structural Equation Modeling, pp. 1–33. Lawrence Erlbaum, Mahaw (2001a)
- Muthén, B.: Second-generation structural equation modeling with a combination of categorical and continuous latent variables: new opportunities for latent class/latent growth modeling. In: Collins, L., Sayer, A. (eds.) New Methods for the Analysis of Change, pp. 291–322. APA, Washington (2001b)
- Muthén, B.: Beyond SEM: general latent variable modeling. *Behaviormetrika* **29**(1), 81–117 (2002)
- Muthén, B.: Statistical and substantive checking in growth mixture modeling: comment on Bauer and Curran (2003). *Psychol. Methods* **8**, 369–377 (2003)
- Muthén, B.: Latent variable analysis: growth mixture modeling and related techniques for longitudinal data. In: Kaplan, D. (ed.) The SAGE Handbook of Quantitative Methodology for the Social Sciences, pp. 345–368. Sage, Thousand Oaks (2004)
- Muthén, B.: Latent variable hybrids: overview of old and new models. In: Hancock, G.R., Samuelsen, K.M. (eds.) Advances in Latent Variable Mixture Models, pp. 1–24. Information Age Publishing, Charlotte (2008)
- Muthén, B., Asparouhov, T.: Growth mixture modeling: analysis with non-Gaussian random effects. In: Fitzmaurice, G., Davidian, M., Verbeke, G., Molenberghs, G. (eds.) Longitudinal Data Analysis, pp. 143–165. Chapman & Hall/CRC Press, Boca Raton (2008)
- Muthén, B., Muthén, L.: Integrating person-centered and variable-centered analysis: growth mixture modeling with latent trajectory classes. *Alcohol. Clin. Exp. Res.* **24**(6), 882–891 (2000a)
- Muthén, B., Muthén, L.: The development of heavy drinking and alcohol-related problems from ages 18 to 37 in a U.S. national sample. *Journal of Studies on Alcohol* **61**, 290–300 (2000b)
- Muthén, L., Muthén, B.: Mplus User's Guide, 4th edn. Muthén & Muthén, Los Angeles (2006)

- Muthén, L., Muthén, B.: Mplus User's Guide, 6th edn. Muthén & Muthén, Los Angeles (2010)
- Muthén, B., Shedden, K.: Finite mixture modeling with mixture outcomes using the EM algorithm. *Biometrics* **55**(2), 463–469 (1999)
- Nagin, D.S.: Analyzing developmental trajectories: a semi-parametric, group-based approach. *Psychol. Methods* **4**, 139–157 (1999)
- Nagin, D.S.: Group-Based Modeling of Development. Harvard University Press, Cambridge (2005)
- Nagin, D.S., Land, K.C.: Age criminal careers, and population heterogeneity: specification and estimation of a nonparametric, mixed Poisson model. *Criminology* **31**(3), 327–362 (1993)
- Nylund, K., Asparouhov, T., Muthén, B.O.: Deciding on the number of classes in latent class analysis and growth mixture modeling. A Monte Carlo simulation study. *Struct. Eq. Model.* **14**, 535–569 (2007)
- Odgers, C.L., Caspi, A., Poulton, R., Harrington, H., Thompson, M., Broadbent, J.M., Dickson, N., Sears, M.R., Hancox, B., Moffit, T.E.: Prediction of adult health burden by conduct problem subtypes in males. *Arch. Gen. Psychiatry* **64**, 476–484 (2007)
- Pöge, A.: Persönliche Codes bei Längsschnittstudien: Ein Erfahrungsbericht. *ZA-Inf.* **56**, 50–69 (2005)
- Rao, C.R.: Some statistical methods for comparison of growth curves. *Biometrics* **14**, 1–17 (1958)
- Reinecke, J.: Delinquenzverläufe im Jugendalter: Empirische Überprüfung von Wachstums- und Mischverteilungsmodellen. Institut für sozialwissenschaftliche Forschung e.V., Münster (2006a)
- Reinecke, J.: Longitudinal analysis of adolescents' deviant and delinquent behavior. Applications of latent class growth curves and growth mixture models. *Methodology* **2**, 100–112 (2006b)
- Reinecke, J.: The development of deviant and delinquent behavior of adolescents. Applications of latent class growth curves and growth mixture models. In: van Montfort, K., Oud, J., Satorra, A. (eds.) *Longitudinal Models in the Behavioral and Related Sciences*, pp. 239–266. Lawrence Erlbaum, Mahwah (2007)
- Reinecke, J.: Klassifikation von Delinquenzverläufen: Eine Anwendung von Mischverteilungsmodellen. In: Reinecke, J., Tarnai, C. (eds.) *Klassifikationsanalysen in Theorie und Praxis*, pp. 189–218. Waxmann, Münster (2008)
- Roeder, K., Lynch, K.G., Nagin, D.S.: Modeling uncertainty in latent class membership: a case study in criminology. *J. Am. Stat. Assoc.* **94**, 766–776 (1999)
- Ross, S.M.: Introduction to Probability Models, 5th edn. Academic Press, New York (1993)
- Rost, J., Langeheine, R.: Applications of Latent Trait and Latent Class Models in the Social Sciences. Waxmann, Münster (1997)
- Schwarz, G.: Estimating the dimension of a model. *Ann. Stat.* **6**(2), 461–464 (1978)
- Tucker, L.R.: Determination of parameters of a functional relation by factor analysis. *Psychometrika* **23**, 19–23 (1958)
- Willet, J.B., Sayer, A.G.: Using covariance structure analysis to detect correlates and predictors of individual change over time. *Psychol. Bull.* **116**(2), 363–381 (1994)