
Question:

WARNING: THE MODEL ESTIMATION HAS REACHED A SADDLE POINT OR A POINT WHERE THE OBSERVED AND THE EXPECTED INFORMATION MATRICES DO NOT MATCH. AN ADJUSTMENT TO THE ESTIMATION OF THE INFORMATION MATRIX HAS BEEN MADE. THE CONDITION NUMBER IS - 0.214D-02. THE PROBLEM MAY ALSO BE RESOLVED BY DECREASING THE VALUE OF THE MCONVERGENCE OR LOGCRITERION OPTIONS OR BY CHANGING THE STARTING VALUES OR BY USING THE MLF ESTIMATOR.

THE MODEL ESTIMATION TERMINATED NORMALLY

And after sending my output you recommended that I could use the results because the output contained standard errors. I followed your advice to ignore the warning. Now me and my supervisors are wondering whether it would not be better to use the same information matrix estimation for all analyses, so is there a way we could make the same adjustment to the information matrix for our other multilevel analyses? We feel they would be better comparable in this way. We have several analyses that have the same input as described above (only different variables) and while several analyses do result in the warning and thus the information matrix is adjusted, others do not, with the result that we now use two different information matrix estimation methods whereas we would like to use the same method for all analyses.

Is this possible in Mplus? Many thanks!

 [Tihomir Asparouhov](#) posted on Wednesday, October 12, 2016 - 4:14 pm

There are 4 suggestions in the last sentence

THE PROBLEM MAY ALSO BE RESOLVED BY DECREASING THE VALUE OF THE MCONVERGENCE OR LOGCRITERION OPTIONS OR BY CHANGING THE STARTING VALUES OR BY USING THE MLF ESTIMATOR.

You should try to use those to resolve the problem before proceeding.

But if the message persists the last suggestion estimator=mlf can be used to avoid the problem and have the same method be used for all SE computations.

Question:

Thank you for your reply. Of course I noticed the 4 suggestions, but when I asked on this forum what the best strategy would be (since I am not a very experienced Mplus user) and sent my output files to the Mplus support desk I was advised: "You receive standard errors. They are in the column S.E. Therefore, you can ignore the message." I also read that the MLF estimator overestimated the standard errors and on the forum I noticed that people were advised against using it in cases comparable to mine. Because (in fact I think it was your article) I also read about the possibility of using a simplified version of the actual information matrix, I figured that this may be another way to solve the problem. For example to assume orthogonality (since there are simulation studies showing that in the type of analyses I am doing this does not have a large effect on the results).

I hope this is a little bit clear! So my question is whether/how I can use a simplified version of the information matrix in my multilevel analyses (for an example of one of my inputfiles see above) and also whether you would recommend this. Or would you rather recommend something else? (If it is changing the starting values after all could you please advise me on which values to choose and how?)

Many thanks in advance!

[Tihomir Asparouhov](#) posted on Thursday, October 13, 2016 - 2:41 pm

There are three things to keep in mind.

- 1) Figure out the reason for the problem
- 2) Decide if you should stick with that model or you should change the model
- 3) If you decide to keep the model what to do to improve the estimation.

For 1) looking at your input this is almost certainly caused by an effect that is not random (near zero variance .. remove it ... use fixed effect) or by two random effects that have a correlation of +1 (I would use estimator=bayes for this problem)

- 2) Given that this is a part of a bigger set of model runs I assume you don't want to change the model but that will be most desirable, however the ML estimator tends to get into singular random effect matrices simply because of small number of clusters and the Bayes estimator switch is best then.
- 3) The saddle point message is 75% of the time incomplete convergence most likely you can resolve using miter=50000; mconv=0.000001; The slow convergence is due to between level singularities. About 25% it is caused by unidentified model.

You best bet apart from that is the MLF estimator. The adjustment amounts to removing negative eigenvalues - not something you can do when there is no problem.

You can add starting values using simpler runs that converge such as models with one or two fewer random effects (I don't think this is likely to solve the problem though) or use starts.

To summarize estimator = bayes or mlf is the best bet or modify the model.

 [Tihomir Asparouhov](#) posted on Thursday, October 13, 2016 - 2:42 pm

One last note - if none of the above improves the situation - it is safe to use the standard errors that Mplus gives you.